



Calling instructions pregnancy

COMMON:

CALL IN THE EVENT OF:

If you have
one or more
of these com-
plaints after a
pregnancy
term of 20
weeks:



- Headache
- Seeing stars/flashes of light
- Vomiting
- Feeling of tightness around the head or upper abdomen
- Pain in the abdomen or between the shoulder blades
- Sudden facial fluid retention
- Tingling in the fingers



Blood loss is always a reason for consultation. A small amount of blood loss can occur during a regular pregnancy term.

Blood loss



- Clear red blood loss

Tip: Do not dispose of your underwear or pantyliner in the event of blood loss



- A regular and identifiable pattern of movements after 28 weeks.

Tip: lying on your left side allows you to feel the movements of the baby best

Baby
movements
after 28 weeks



- Less baby movements than you normally experience
- During relaxation and paying attention, there is less than 10 times movements within 2 hours
- Concern about the amount or strength of the movements



- Urine
- Discharge, also when more watery than normal

Fluid loss
before 37
weeks



- Constantly wet underwear
- Continuous small doses of fluid loss
- Green/brown fluid loss



- Prelabour when active or when the baby is moving
- Not painful
- No regular pattern

Prelabour/
cramps before
37 weeks



- Worsening of cramps
- There is regularity (every half hour or more frequent)
- Painful

Concern



- Always call if concerned or worried
- Any questions on the information supplied



CALL YOUR MIDWIFE CARE WORKER ON THE EMERGENCY NUMBER